

ENROFLOX INJECTION  
FOR DOGS 2.27%  
20 mL

ENROFLOX INJECTION  
FOR DOGS 2.27%  
50 mL

ENROFLOX INJECTION  
FOR DOGS 2.27%  
100 mL

▼ FOLD HERE ▼

## Enroflox<sup>®</sup> (enrofloxacin) Injection for Dogs 2.27% Dosing



Enroflox<sup>®</sup> Injection may be used as the initial dose at 2.5 mg/kg. It should be administered intramuscularly (IM) as a single dose followed 12 hours later by initiation of Enroflox Chewable Tablets.



WEIGHT	ENROFLOX <sup>®</sup> INJECTION FOR DOGS 2.5 mg/kg
10 lb (4.5 kg)	0.5 mL
15 lb (6.8 kg)	0.75 mL
20 lb (9.1 kg)	1.00 mL
25 lb (11.3 kg)	1.25 mL
30 lb (13.6 kg)	1.50 mL
35 lb (15.9 kg)	1.75 mL
40 lb (18.1 kg)	2.00 mL
45 lb (20.4 kg)	2.25 mL
50 lb (22.7 kg)	2.50 mL

**CAUTION:** Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extralabel use of this drug in food-producing animals. **CONTRAINDICATIONS:** Enrofloxacin is contraindicated in dogs known to be hypersensitive to quinolones. The safe use of enrofloxacin has not been established in large and giant breeds during the rapid growth phase. The use of enrofloxacin is contraindicated in small and medium breed dogs during the rapid growth phase (between 2 and 8 months of age).

**WARNINGS:** For use in animals only. The use of this product in cats may result in Retinal Toxicity. Keep out of reach of children. Observe label directions and see product labeling for full product information.

© 2022. All rights reserved. The Norbrook logos and Enroflox are registered trademarks of Norbrook Laboratories Limited.

0422-ENR-M02C