

100 x 4g

SureSeal

READ THE ENTIRE LABEL BEFORE USE

PRESENTATION:

SURESEAL is a sterile intramammary suspension. Each syringe contains 4.0 g of suspension incorporating 2.6 g of Bismuth subnitrate. Supplied in outers of 24, 60, 100 and 120 unit-dose syringes.

INDICATIONS:

SURESEAL is indicated for the prevention of mastitis during the non-lactating (dry) period:

- in lactating heifers (first calves) and cows with somatic cell counts (SCC) of less than 120,000 and 150,000 respectively.
- in cows and heifers with a higher somatic cell counts (SCC), or immediately following treatment with Noroclox DC 600 (A9281), Cloxamp DC 500 (A8038), Cloxamp DC 600 (A8039) or Durodry (A7862). Refer also to the labels for these products.

Sureseal is also indicated for the prevention of peri-calving mastitis in maiden heifers.

DIRECTIONS FOR USE:

Pinch the top of the teat and infuse the sealant, ensuring that only the teat is filled. When the syringe is nearly empty, withdraw the syringe and deposit a small amount in the teat canal. If teats are small, infuse as normal but discard any remaining sealant that is not needed to fill the teat. Do not reuse leftover sealant.

DOSAGE & ADMINISTRATION:

Lactating heifers and cows with SCC below 120,000 and 150,000 respectively:

Immediately after the last milking at drying off, prior to treatment with SURESEAL, the udder should be stripped.

The recommended method is to disinfect and treat one teat at a time. Clean teat thoroughly with a teat wipe (or cotton wool ball) moistened in 70% methylated spirits (30% water), and immediately infuse the entire contents of one SURESEAL syringe to form a seal. Repeat the process on the other three teats and finish by teat spraying all teats. After infusion do not massage or squeeze the teat, as this may dislodge the seal.

Lactating heifers and cows with SCC above 120,000 and 150,000 respectively:

Immediately after the last milking at drying off, prior to treatment with dry cow therapy and SURESEAL, the udder should be stripped. The recommended method is to disinfect and treat one teat at a time. Clean teat thoroughly with a teat wipe (or cotton wool ball) moistened in 70% methylated spirits (30% water). Insert Noroclox DC 600 (A9281), Cloxamp DC 500 (A8038), Cloxamp DC 600 (A8039) or Durodry (A7862), and massage into the quarter as per the respective product label. Redisinfect the teat. Carefully infuse the entire contents of one SURESEAL syringe to form a seal. Repeat the process on the other three teats and finish by teat spraying all teats. After infusion do not massage or squeeze the teat, as this may dislodge the seal.

Maiden heifers:

Sureseal should only be administered to maiden heifers in herds where heifer mastitis levels are clinically significant. Care should be taken to ensure administration of Sureseal in heifers is performed to best practice recommendations. Administer Sureseal pre-calving before springing ('bagging up') of the udder; ie from approximately four weeks to one week prior to the planned start of calving in a typical seasonally calving herd. It is not recommended to administer Sureseal closer than one week prior to calving.

REMOVAL

It is important to remove SURESEAL from fresh cows to stop any particles entering the vat.

Thoroughly hand strip 10-12 strips per quarter at the first milking after calving. Stripping should commence at the very top of the teat. Where possible, avoid removal of SURESEAL by the milking machine.

CONTRA-INDICATIONS:

Do not use in lactating cows.
Do not use in cows with clinical mastitis.

WITHHOLDING PERIODS:

MILK: Milk intended for sale for human consumption must be discarded during treatment for not less than **8 milkings** or approximately **96 hours** after calving.

MEAT: Nil

Handling Precautions:

Wear protective gloves when handling this product.

First Aid:

For advice contact the National Poisons Centre 0800 POISON (0800 764 766). Get medical attention if you feel unwell. Have this product label with you when you call.

Disposal:

Dispose of product, packaging and waste at an approved landfill or other equivalent facility.

WARNINGS:

Do not warm the product before use.
To ensure maximum efficacy of the product, best drying off practice according to national guidelines should be followed both before and after drying off animals when using Sureseal. It is important that infection of the udder does not occur when administering any dry cow products. Accidental introduction of bacteria into the quarter during insertion of the intramammary syringe can result in serious mastitis.

PHARMACEUTICAL PRECAUTIONS:

Partly used syringes must be discarded.
Store below 25 °C.
KEEP OUT OF REACH OF CHILDREN.



Registered pursuant to the ACVM Act 1997 No. A10866
See www.foodsafety.govt.nz for registration conditions.

Registered to and Distributed by:

Norbroke NZ Ltd.,
KPMG Centre,
18 Viaduct Harbour Avenue,
Auckland.
Ph: 0800 224 022

Manufactured by:

Norbroke Laboratories Ltd.,
Station Works,
Newry,
Co. Down,
Northern Ireland.

Batch:

Expiry:

062460L05



Norbroke